




1/29



2/29




3/29




4/29

FIG. 4




5/29

FIG. 5

6/29

FIG. 6

Preparation of HVJ envelope vector



7/29

FIG. 7

Protein profiles of HVJ, UV-inactivated HVJ,
and HVJ envelope vector




FIG. 8

8/29

Electron micrograph of an HVJ envelope vector

(1) Untreated HVJ



100 nm

(2) HVJ containing no DNA, which was subjected to an octylglucoside treatment



(3) HVJ containing DNA, which was subjected to an octylglucoside treatment



09937839-02102

09/937839

FIG. 9A

9/29

Effects of octylglucoside on gene transfer
by HVJ envelope vector






FIG. 9B



10/29

FIG. 9C

Effects of octylglucoside on gene transfer
by HVJ envelope vector




201210 6632550

FIG. 10A

11/29


Effects of protamine sulfate on gene transfer by HVJ envelope vector



12/29

FIG. 10B



Effects of protamine sulfate on gene transfer by HVJ envelope vector



13/29

FIG. 11A


Effects of DNA amounts on gene expression using frozen HVJ envelope which has been treated with octylglucoside



14/29

FIG. 11B


Effects of DNA amounts on gene expression
by HVJ envelope vector



201220" 66346550

FIG. 12

Effects of HVJ titer on gene expression




09/937839


16/29

FIG. 13A



Study of irradiation amount in UV inactivation

**FIG. 13B**

Study of treatment concentration in BPL inactivation




17/29

FIG. 14**SAS: Effects of infection time and protamine sulfate concentration****FIG. 15****HAEC: Effects of infection time and protamine sulfate concentration**

18/29

FIG. 16A

Luciferase activity by HVJ envelope —
vector in mouse liver




201220"6E84E660

19/29

FIG. 16B

Luciferase activity by HVJ envelope vector
in mouse uterus



09/937839

20/29

201220" 6EB/E66B

FIG. 16C

LacZ expression by HVJ envelope vector in mouse uterus



21/29

FIG. 16D

Gene transfer into central nervous system using new HVJ

#1 HVJ-GFP

HVJ-GFP of 10,000 HAU was administered to SD rats (male, body weight: 300 to 400 g) via the cisterna magna or via the carotid artery. Samples were taken three to four days later.

Live sections were prepared, which were subjected to observation under fluorescence microscopy.

(administration via the cisterna magna) ①

Incorporation into the brain surface was confirmed.


No incorporation into deep portions of the brain was confirmed.

No incorporation into the choroid plexus was confirmed. → administration via the cisterna magna is considered to result in permeation through the intrathecal space, so that expression is usually observed in the choroid plexus


(administration via the carotid artery) ②, ③

Significant expression was confirmed on the administered side (left side). Expression was confirmed not only in the brain surface portions but also in the basal ganglia portion. Expression was also confirmed in the brain surface of the other brain, which was considered to have resulted from a flow to the other side through a collateral flow.

① Administration via the cisterna magna



② Administration via the carotid artery



③ Administration via the carotid artery




09/937839

22/29

201220" 6E8/660

FIG. 16E Inhibition of VEGF-induced angiogenesis
by gene transfer using HVJ



09/937839

23/29

201220" 6E84E660

FIG. 16F



24/29


20F220" 6E84E660

Introduction of FITC-ODN into BHK-21 cells by HVJ envelope vector


FIG. 17A

60 min


Fluorescence image



Phase-contrast image


**FIG. 17B**

10 min



25/29

FIG. 18A Gene transfer into NALM-6 by centrifugation with HVJ envelope




Protamine sulfate concentration and centrifugation

201220 668/E660

09/937839

26/29

FIG. 18BGene transfer into CCRF-CEM by:
centrifugation with HVJ envelope

Protamine sulfate concentration and centrifugation


09/937839

27/29

FIG. 18C

Gene transfer into K-562 by centrifugation
with HVJ envelope

(15 K rpm, 10 min, 20°C)




201220" 6E8/E660

09/937839

28/29

FIG. 19

Gene transfer into mouse melanoma (B16-F1)
mass using HVJ envelope



201220-6834650

09/937839

29/29

FIG. 20




FIG. 21

